

Remarks/Arguments

Applicants respond to Examiner's objections as follows in accordance with the listing on pages 3 -6 of the Office Action:


- A. Amended in the attached substitute specification per Examiner's request.
- B. Amended in the attached substitute specification per Examiner's request.
- C. Amended in the attached substitute specification per Examiner's request.
- D. Amended in the attached substitute specification per Examiner's request.
- E. Amended in the attached substitute specification per Examiner's request.
- F. Amended in the attached substitute specification per Examiner's request.
- G. Amended in the attached substitute specification per Examiner's request.
- H. Applicant clarifies that it is not usual practice to provide the stem length because it is not an accurate measurement as it varies based on environmental and cultural conditions such as age and juvenility or maturity of the plant. The internode length, which is included in the specification, is a more accurate measurement. No amendment has been made to the specification.
- I. Amended in the attached substitute specification per Examiner's request.
- J. Amended in the attached substitute specification per Examiner's request.
- K. Amended in the attached substitute specification per Examiner's request.
- L. Amended in the attached substitute specification per Examiner's request.
- M. Amended in the attached substitute specification per Examiner's request.
- N. The plant shown in the photographs is not the plant described in the specification; however, it is of the same cultivar. A clarifying amendment has been made in the attached substitute specification per Examiner's request.

CONCLUSION

Applicant submits that in view of the foregoing remarks and/or amendments, the application is in condition for allowance, and favorable action is respectfully requested. The Commissioner is hereby authorized to charge any fees, including extension fees, which may be required, or credit any overpayments, to Deposit Account No. 50-1001.

Respectfully submitted,

Date: July 14, 2004


Bradley Ganz
Registration No. 34,170
P. O. Box 10105
Portland, Oregon 97296
Telephone: (503) 224-2713
Facsimile: (503) 298-2172
email: mail@ganzlaw.com

Substitute Specification - Marked-Up

Impatiens Plant Named 'TILav Improved'**Latin Name of the Genus and Species and Variety Denomination**

- 5 *Impatiens walleriana* plant and known by the cultivar name ~~'TILav Improved'~~ 'TILav Improved'.

BACKGROUND OF THE INVENTION**Field of Invention**

- 10 The present invention comprises a new and distinct cultivar of plant, botanically known as *Impatiens walleriana* and hereinafter referred to by the cultivar name of 'TILav Improved'.

1. The plants are useful in landscaping, window and hanging baskets and flower gardens.

- 15 2. The plant is an *Impatiens walleriana*.

3. The plants are generally classed as an annual bedding plant.

Description of the Relevant Prior Art

- 20 1. The pollen parent is an unpatented pollen producing double *Impatiens walleriana* coded P-BR-131, and the seed parent is an unpatented semi-double *Impatiens* coded M-BR-203. 'TILav Improved' is both male and female sterile. Cultivar is both female and male sterile. Both seed parents are capable of both seed and pollen production.

Substitute Specification – Marked-Up

2. The seed parent was a Pink flowered cultivar and the pollen parent was a Lavender flowered cultivar, neither parent is either patented or commercially available. The cross was made in the breeder's controlled breeding program at Rogue River, Oregon by lateral stems with two or more leaves. The first asexual reproduction was also made at Rogue River, Oregon. Successive asexual generations produced from cuttings with two or more leaves have shown the cultivar to be stable and reproduced true to type.

COMPARISON

1. The plant is compared to 'TiLav', [US PP] U.S. Plant Patent 13,003.
2. The flower color of 'TiLav Improved' is lighter than that of 'TiLav'.
3. The Foliage color of 'TiLav Improved' is lighter than that of 'TiLav'.
4. 'TiLav Improved' flowers better in temperatures above 90° F than 'TiLav'.
5. See comparison chart labeled Chart 1.

Summary of the Invention

The cultivar is unique either in one or a combination of the characteristics described herein. It is a new, double *Impatiens walleriana* producing lavender flowers; dark green foliage; medium-vigorous growth habit; fully double flowers held above or beyond the foliage on strong peduncles and pedicels; a compact mounded growth habit.

Substitute Specification – Marked-Up

DESCRIPTION OF THE PHOTOGRAPHS

The attached color photographs illustrate a plant of 'TiLav Improved' grown in winter and just beginning to mature. The plant shown in the photographs is not the plant described in the specification; however, it is of the same cultivar. Variations between the written description and the colors in the photographs may occur due to the type of film and/or light reflectance and the type of light captured by the film. In the event there are such variations, the written description shall control.

PLANT DESCRIPTION

The following description was taken at Rogue River, Oregon on ~~February 7, 2003~~ February 7, 2004. The description is of a plant of 'TiLav Improved' similar to the one shown in the photograph.

The following description was taken from a plant approximately 12-weeks of age under the following environment and cultural practices at Rogue River, Oregon. Phenotypical variations may result in plants grown under conditions and locations other than those listed herein.

Container – 6 inch azalea pot.

Medium – Peat-lite type of soilless medium.

Greenhouse covering – Double layer Polyethelene with 50% shade applied above the covering.

Minimum temperature – 65° F.

Maximum temperature – 75° F.

Light levels – 1000 to 1500 ft candles.

Fertilizer – 20-10-20 with trace elements applied twice followed by one leaching with clear water.

3

Ganz Law, PC
P.O. Box 10105
Portland, Oregon 97298
Phone: (503) 224-2713
DocId: No. GOS-2-089
Express Mail No: EV328263891US

Substitute Specification – Marked-Up

Cutting type – Lateral stems with two or more leaves.

Propagation:

Type of Cutting: Lateral stems with two or more leaves.

5 *Time to Initiate Roots:* Generally about 7 to 10 days at soil temperature of 72° F.

Appearance and Form of Mature Plant:

Note: All size measurements for plant, stems, foliage, flowers, peduncles, pedicels, buds, and sepals are taken during winter, low light conditions when the above are the smallest. Spring and summer high light sizes are generally much larger than the sizes that are described
10 below.

Growth habit: Mounded and of compact-medium vigor.

Plant Size: Plants mature at about 20 cm in height and about 35 cm in width. Both of these measurements are a function of age, environmental and cultural practices; and can vary accordingly.

15 *Rooting Habit:* The roots are fibrous and well branched.

Rooting Ability: Easy, no hormones needed.

Stems: Stems are freely produced, and no pinching to induce branching is needed. Color is translucent 146C to 146D with stippling on juvenile stems of very small markings that are difficult to determine but which appear to be close to 178A. The stippling is heaviest at the
20 nodes. The stippling appears to be outgrown as mature stem color becomes solid 146C to 146D. The stems at maturity are generally at least 3 mm in diameter at the internodes, and the internode length is generally at least 0.5 cm in length.

Foliage: Shape is ovate with cuneate to attenuate base; crenate margin, margin having tiny spines of an undeterminable greenish color, and each being less than 1 mm in length and

Substitute Specification – Marked-Up

protruding outwards in a perpendicular direction from leaf edge and parallel to the flat surfaces of the leaf, between the lobes of the crenate margin; and an acuminate apex. Size of largest is about 3 cm long; about 2.75 cm wide.

Foliage Color: Adaxial surface close to 146A; abaxial surface close to 148D; adaxial surface venation is close to 146D at base, becoming barely distinguishable to undistinguishable towards the leaf margins; abaxial surface venation, main vein at base is closest to 146A to 146B, lateral veins are close to 146A.

Foliage Texture: Smooth and flat.

Petioles: Longest petioles are about 2.5 cm long, half-round, flat to slightly convex on upper surface, which is about 2 mm wide, depth of about 1.5 mm. Adaxial surface color appears to be closest to 146C; abaxial surface color appears to be close to 146C to 146D.

Flower Habit: Flowers are produced continuously throughout the flowering season.

Natural Flowering Season: All year in greenhouse conditions, and the frost-free period from spring through fall, when planted outdoors.

Duration of flower: Each flower generally lasts about 5 days at 75° F maximum and 70° F minimum temperature.

Flowers Borne: Above or beyond the foliage.

Flower Texture: Smooth and satiny.

Flower Color: Petals/petaloids - Adaxial surface is close to 75A. As the flower ages, the color at the edges fades toward 75B, and fades more towards the middle, changing to 69A to 69B. There is a basal spot that varies in intensity of color, generally being either close to 67A, or is more intense and close to 59B to 59C. The petaloids toward the center of the flower tend to have the basal spot color closest to 67A, while 59B to 59C tends to be more common on the

Substitute Specification – Marked-Up

outer petal's basal spot. Abaxial surface is closest to 75D, there is generally but not always a basal spot close to 67A.

Quantity. Usually two or more open flowers per stem, and generally 15 or more from visible buds to fully mature flowers.

5 *Petal/Petaloid count:* Generally 25 or more.

Size: ~~Flowers are~~ Generally, generally at least 3.5cm [wide] in diameter; generally at least 1 cm deep with largest petals/petaloids. The largest petals/petaloids are generally at least 1.5 cm wide and at least 1.5 cm [deep] long; obovate to exaggerated obovate in shape with obtuse to retuse apex, entire margin, and cuneate to obtuse base. The above
10 measurements refer to the average flowers[;]—

Flower Buds: The buds before opening are about 1 cm in length; about 0.75 cm in diameter; shape is ovate. Bud color prior to opening is close to 145A on top and close to 145D on bottom.

Peduncles: Generally at least 1.25 cm long; about 2 mm in diameter; colored close to
15 144B. Good strength.

Pedicels: Generally two or more per peduncle. Each is generally at least 1.5 cm in length, and at least 1.5 mm in diameter. Color is close to 144B. Good strength.

Calyx: Calyx consists of generally one sepal and one spur. The spur is generally at least 1.75 cm long, and about 2 mm in diameter at sepal end; shape is an acicular tapering tube
20 about 1 mm in diameter at the apex, and it is attached toward the base of the sepal in a peltate manner. Color is close to 178A at apex, is an orangish-green that neither matches, nor is very close to any color in the chart, with the closest color in the chart being 166D at base. The apical color generally changes with maturity to close to 160A.

Substitute Specification – Marked-Up

- The sepal generally measures at least 0.75 cm wide and at least 1.1 cm long; ovate in shape with an acute apex, entire margin, and obtuse to cordate base; adaxial surface is lighter than 145D, with a base spot of close to 74A to 74B where the spur connects; the abaxial surface is close to 145D. The adaxial basal spot sometimes appears through to the abaxial side, giving an apparent spot that appears close to 70A where the spur attaches.

Reproductive Organs: The reproductive organs are replaced with sterile petaloids.

Disease Resistance: Disease resistance has not been tested.

Dampness Resistance: The plant has shown a good ability to hold the flowers in an outward manner during summer rains

- Heat Tolerance:** The plant has shown a good tolerance to temperatures exceeding 100° F at Rogue River by flowering continuously during temperatures as high as 113° F.

CHART 1 – COMPARISON CHART TO PRIOR ART

15

	'TILav Improved'	'TILav'
Stems – color	Color is translucent 146C to 146D with stippling on juvenile stems that is heaviest at the nodes which appear to be close to 178A. The stippling appears to be outgrown on mature stems as the stem color becomes solid 146C to 146D	147A, spots of 187A
Foliage – Size	Size of largest is about 3 cm long; about 2.75 cm wide.	
Abaxial color	148D	148B, with marking close to 177A.
Foliage – Abaxial venation color	Main vein at base is closest to 148A to 146B, lateral veins are close to 146A.	148A

7

Ganz Law, PC
P.O. Box 10105
Portland, Oregon 97206
Phone: (503) 224-2713
Docket No.: 009-2-009
Express Mail No: EV329253891US

Substitute Specification – Marked-Up

Foliage- Adaxial color	Close to 146A	147A
Foliage – Adaxial venation color	Close to 146D at base becoming barely distinguishable to undistinguishable towards the leaf margins;	147A
Petiole- Size	Longest petioles about 2.5 cm long, about 2 mm wide, depth of about 1.5 mm.	
Adaxial color	Closest to 146C.	146B, 178A
Petiole Abaxial color	Close to 146C to 146D	146B
Petal/Petaloid –Size	Generally at least 1.5 cm wide and at least 1.5 cm [deep] long.	2 cm long, 2.5 cm wide.
Abaxial color	Closest to 75D, there is generally but not always a basal spot of close to 67A.	75A
Petal/Petaloid Adaxial color	Close to 75A, as the flower ages, the color at the edges fades toward 75B and fades more towards the middle changing to 69A to 69B, there is a basal spot that varies in intensity of color generally being either close to 67A or is more intense and being close to 59B to 59C. The petaloids toward the center of the flower tend to have the basal spot color closest to 67A while 59B to 59C tends to be more common on the outer petals basal spot.	Mature flowers 74D with dark basal spot of 59B; Juvenile color 74C with basal spot of 59B.
Peduncle – Size	Generally at least 1.25 cm long; about 2 mm in diameter;	2 cm long; 2.5 mm diameter.
Color	Close to 144B.	148A
Pedicel- Size	Generally at least 1.5 cm in length, at least 1.5 mm in diameter.	2 cm long; 1.5 mm diameter.
Color	Close to 144B.	146B
Sepal –Size	Generally at least 0.75cm	1 cm long; 0.8 cm wide.

Substitute Specification – Marked-Up

Adaxial color	wide, and at least 1.1 cm long; Lighter than 145D, with a base spot of close to 74A to 74B where the spur connects.	145B to 145C with basal spot close to 183C.
Sepal Abaxial color	Close to 145D sometimes an apparent spot that appears close to 70A where the spur attaches.	145C
Spur- Size	Generally at least 1.75 cm long, and about 2mm in diameter at sepal end, about 1 mm in diameter at the apex.	3cm long, 2 mm wide.
Color	Close to 178A at apex, is an orangish-green that neither matches, nor is very close to any color in the chart with the closest color in the chart being 168D at base. The apical color generally changes with maturity to close to 160A.	146C at base, 183B at apex.
Buds- Size	About 1 cm. in length; about 0.75 cm. in diameter.	0.8 cm long; 0.6 cm wide; 0.7 cm depth;
Color	Close to 145A on top and close to 145D on bottom.	146B on top; 146D on bottom.

Substitute Specification – Marked-Up

CLAIM

IT IS CLAIMED:

A new and distinct cultivar of double *impatiens walleriana* plant named 'TiLav Improved',
as illustrated and described.

5

10

10

Ganz Law, PC
P.O. Box 10105
Portland, Oregon 97206
Phone: (503) 224-2713
Docket No.: COS-2.009
Express Mail No: EV329253691US

Substitute Specification -- Marked-Up

ABSTRACT

A new double *impatiens walleriana* plant named 'TiLav Improved'; producing lavender flowers; dark green foliage; medium-vigorous growth habit; fully double flowers held above or
5 beyond the foliage on strong peduncles and pedicels; and a compact mounded growth habit.

Substitute Specification – Marked-Up

COLOR CODE CHART

	Stems:	146C, 146D, 178A.
	Foliage Abaxial surface:	148D
	Foliage Abaxial surface venation:	146A, 146B
5	Foliage Adaxial surface:	146A
	Foliage Adaxial surface venation:	148D
	Petiole Abaxial surface:	146C
	Petiole Adaxial surface:	146C, 146D
	Petals/Petaloids Abaxial surface:	75D, 67A
10	Petals/Petaloids Adaxial surface:	75A, 75B, 69A, 69B, 67A, 59B, 59C, 67A, 59B, 59C
	Buds:	145A, 145D
	Peduncles:	144B
	Pedicels:	144B
	Sepals adaxial surface:	145D, 74A, 74B
15	Sepals abaxial surface:	145D, 70A
<hr/>		
	Spur:	178A, 166D, 160A